


HIV AMONG MEN IN MARYLAND, 2013


MAY 2015

New HIV Diagnoses

- Men made up 71.0% (990) of the 1,395 reported adult/adolescent HIV diagnoses (age 13+) in Maryland in 2013.
- The proportion of HIV diagnoses in Maryland has decreased in men from 85.2% (in 1985) to 71.0% (in 2013).
- 71.5% of the 990 reported HIV diagnoses among men (age 13+) in 2013 were among non-Hispanic (NH) blacks, 17.2% NH white, 6.3% Hispanic, and 5.1% other races.
- Most HIV diagnoses in men in 2013 were among Men who have Sex with Men (MSM) (69.3%).



- Total living HIV case rates males (all ages) were
 4-8 times higher among NH blacks (1,781.9 cases per 100,000) as compared to other races.
 In other words, 1 in every 56 NH black males in Maryland was reported to be living with HIV.
- Of the 19,667 total living HIV cases among males in Maryland, about half (50.5%) had a viral load test result reported in 2013. Among the 9,926 with a reported test result, 72.4% had a suppressed viral load.



Characteristics of Men Living with HIV


Exposure Category

- 50.4% MSM
- 25.9% IDU
- 18.3% HET
- 5.0% MSM/IDU

Ť

Race/Ethnicity

- 72.2% NH Black
- 17.8% NH White
- 5.6% Hispanic
- 4.3% NH Other
- Of the 30,306 total living HIV cases (age 13+) reported in Maryland in 2013, 64.9% (19,667) of those were among males. Adult/adolescent living case rates (per 100,000) were twice as high in men (825.5) as compared to women (412.2). In other words, 1 in 121 men was reported to be living with HIV in Maryland.



MSM: Men who have Sex with Men | HET: Heterosexual Exposure | IDU: Injection Drug User